GET THE FACTS

Available Outpatient COVID-19 Treatments

Sick with COVID-19 symptoms?

You may be eligible for treatments that are now available for some non-hospitalized adults and children with COVID-19.

These prescription treatments:

- ✓ are FDA approved or authorized.
- decrease serious disease and hospitalizations.
- work best when used early.
- largely remain effective against recent omicron strains (including XBB.)

For use within 5 days of first symptoms:

NIRMATRELVIR/RITONAVIR (PAXLOVIDTM)

2-3 pills twice a day for 5 days



2x for



MOLNUPIRAVIR (LAGEVRIO®)

4 pills twice a day for 5 days







For use within 7 days of first symptoms:

REMDESIVIR (VEKLURY®)

1 IV infusion (i.e., delivered through the vein) per day for 3 days







For use within 8 days of first symptoms:

in people who are immunocompromised or taking immune-suppressing medicines

HIGH-TITER CONVALESCENT PLASMA

Single IV transfusion over 60 minutes



1x for



Good to Know

4

There is only a short time window when these treatments can be used. So at first sign of illness, get tested for COVID-19 and contact your health care provider or visit covid.gov.



There may be issues with drug interactions using nirmatrelvir/ritonavir (Paxlovid™) with other common medicines, so check with your health care providers.



Molnupiravir should NOT be given to children or pregnant people, and:

- Women should use contraception during the course.
- Men should use contraception during the course and for 3 months after.



These medications are not a substitute for vaccination or other methods to prevent COVID-19.

COVID-19 Real-Time Learning Network

Brought to you by CDC and Rail In SA



For more information on therapeutics, visit: **COVID19LearningNetwork.org**









